



Aléatico N



Name of vine variety in France

Aléatico

Origin

This variety has been found in Corsica for a long time and would seemingly be from Italy (Tuscany).

Synonymy

There is no officially recognized synonym in France, nor in other countries in the European Union.

Regulations

In France, Aléatico N is officially listed in the "Catalogue of vine varieties". This variety is likewise listed in the catalogues of other European Union countries including: Italy and Malta.

Use

Wine vine variety

Evolution of area under vines in France

| | 1958 | 1988 | 1998 | 2008 | 2016 |
|----|------|------|------|------|------|
| ha | 42 | 25 | 30 | 14 | 20 |

Description

Identification signs include:

- the tip of the young shoot has a sparse to moderate coat of flat-lying hairs,
- the young leaves are reddish and bronze color,
- adult leaves are 3 or 5-lobed, with a slightly open petiolar sinus, long teeth with straight sides, an absence of anthocyanin pigmentation and the underside with an absence or very sparse coat of upright hairs and a very sparse coat of flat-lying hairs,
- the berries are round or slightly flattened, slightly muscat flavor with a bluish black skin when ripe.

Genetic profile

| Microsatellite | VVS2 | VVMD5 | VVMD7 | VVMD27 | VRZAG62 | VRZAG79 | VVMD25 | VVMD28 | VVMD32 |
|----------------|------|-------|-------|--------|---------|---------|--------|--------|--------|
| Allel 1 | 131 | 223 | 239 | 176 | 186 | 250 | 248 | 235 | 263 |
| Allel 2 | 133 | 225 | 249 | 191 | 196 | 256 | 254 | 245 | 271 |

Phenology

Bud burst: 2 days after Chasselas. Grape maturity: period II, 2 weeks after Chasselas.

Suitability for cultivation and agronomic production

This variety is fairly fertile and has good vigor. It has upright growth and can be trained and short pruned. It is recommended for hot and dry land.

Sensitivity to diseases and pests

Aléatico N is very sensitive to powdery mildew, grey rot and acid rot.

Technological potential

The clusters and berries are moderate size. This variety with its muscat aromas has good sugar accumulation in the berries and satisfactory total acidity. It makes wines with a lively, shiny but not very intense color. It would appear well adapted for making natural sweet wines. the muscat flavor is oftentimes slight and sometimes fleeting.





Clonal selection in France

The 7 approved Aléatico N clones carry the numbers 802, 859, 860, 861, 892, 893 and 894.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi.
- Documentation interne du Domaine de Vassal. 1949-2011, INRA, Marseillan-plage.
- Cépages et vignobles de France, tome 2. P. Galet, 1990, Ed. Dehan, Montpellier.
- Traité général de viticulture, Ampélographie. P. Viala et V. Vermorel, 1901-1910, Ed. Masson, Paris.

Description of clones approved in France

| Clone number | Identity and availability | | Agronomic data | | Technical data | |
|---|-----------------------------|-----------------------------|--------------------------------|--------------------------------|---------------------------|------------------------------|
| | <i>Origin</i> | <i>Selection</i> | <i>Fertility</i> | <i>Production level</i> | <i>Sugar content</i> | <i>Potential color</i> |
| | <i>Year approved</i> | <i>Agronomic references</i> | <i>Weight of grape bunches</i> | <i>Vigor</i> | <i>Total acidity</i> | <i>Tannic structure</i> |
| | <i>Growing surface area</i> | | <i>Size of berries</i> | <i>Sensitivity to Botrytis</i> | <i>Aromatic intensity</i> | <i>Oenological aptitudes</i> |
| 802 | Corse-du-Sud | ENTAV | medium | high | medium | |
| | 1985 | Corse-du-Sud | high | medium | | |
| | | | medium | | | |
| ENTAV  | | | | | | |
| 859 | Corse-du-Sud | ENTAV | | | | |
| | 1986 | Corse-du-Sud | | | | |
| | | | | | | |
| ENTAV  | | | | | | |
| Small distribution clone conforms with vine variety characteristics. Study underway. | | | | | | |
| 860 | Corse-du-Sud | ENTAV | | | | |
| | 1986 | Corse-du-Sud | | | | |
| | | | | | | |
| ENTAV  | | | | | | |
| Small distribution clone conforms with vine variety characteristics. Study underway. | | | | | | |
| 861 | Corse-du-Sud | ENTAV | | | | |
| | 1986 | Corse-du-Sud | | | | |
| | | | | | | |
| ENTAV  | | | | | | |
| Small distribution clone conforms with vine variety characteristics. Study underway. | | | | | | |
| 892 | Corse-du-Sud | ENTAV | | | | |
| | 1987 | Corse-du-Sud | | | | |
| | | | | | | |

| Clone number | Identity and availability | | Agronomic data | | Technical data | |
|--------------|-----------------------------|-----------------------------|--------------------------------|--------------------------------|---------------------------|------------------------------|
| | <i>Origin</i> | <i>Selection</i> | <i>Fertility</i> | <i>Production level</i> | <i>Sugar content</i> | <i>Potential color</i> |
| | <i>Year approved</i> | <i>Agronomic references</i> | <i>Weight of grape bunches</i> | <i>Vigor</i> | <i>Total acidity</i> | <i>Tannic structure</i> |
| | <i>Growing surface area</i> | | <i>Size of berries</i> | <i>Sensitivity to Botrytis</i> | <i>Aromatic intensity</i> | <i>Oenological aptitudes</i> |

ENTAV  INRA®

Small distribution clone conforms with vine variety characteristics. Study underway.

| | | | | | | |
|-----|--------------|--------------|--|--|--|--|
| 893 | Corse-du-Sud | ENTAV | | | | |
| | 1987 | Corse-du-Sud | | | | |
| | | | | | | |

ENTAV  INRA®

Small distribution clone conforms with vine variety characteristics. Study underway.

| | | | | | | |
|-----|--------------|--------------|--|--|--|--|
| 894 | Corse-du-Sud | ENTAV | | | | |
| | 1987 | Corse-du-Sud | | | | |
| | | | | | | |

ENTAV  INRA®

Small distribution clone conforms with vine variety characteristics. Study underway.



Cette œuvre est mise à disposition selon les termes de la [Licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Partage dans les Mêmes Conditions 4.0 International](https://creativecommons.org/licenses/by-nc-sa/4.0/)

