



Rivairenc blanc B



Name of vine variety in France

Rivairenc blanc

Origin

This vine variety corresponds to the white mutation of Rivairenc N.

Synonymy

In France, this vine variety can officially be called "Aspiran blanc B" for plant propagating material.

Regulations

In France, Rivairenc blanc B is officially listed in the "Catalogue of vine varieties".

Use

Wine vine variety

Evolution of area under vines in France

	1958	1979	2016
ha	2	1	1

Description

The description corresponds to Rivairenc N, except that the berries when ripe have a greenish-yellow skin color.

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allel 1	131	223	247	176	204	252	240	227	239
Allel 2	131	223	249	178	204	260	248	257	261

Phenology

Bud burst: 9 days after Chasselas.
Grape maturity: period II, 3 and 1/2 weeks after Chasselas.

Suitability for cultivation and agronomic production

Characteristics are similar to Rivairenc N.

Sensitivity to diseases and pests

The behavior of Rivairenc blanc B to diseases is comparable to Rivairenc N.

Technological potential

Grape clusters and berries are large. Rivairenc blanc B produces light and neutral wines with low sugar potential.

Clonal selection in France

There are no approved clones for this vine variety at the present time.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi.
- Documentation interne du Domaine de Vassal. 1949-2011, INRA, Marseillan-plage.
- Cépages et vignobles de France, tome 1. P. Galet, 1990, Ed. Dehan, Montpellier.
- Ampélographie. P. Viala et V. Vermorel, 1902-1910, Ed. Masson, Paris.



Cette œuvre est mise à disposition selon les termes de la [Licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Partage dans les Mêmes Conditions 4.0 International](https://creativecommons.org/licenses/by-nc-sa/4.0/)



INRA
SCIENCE & IMPACT



SupAgro
Montpellier

GenoVigne



Pl@ntNet



agropolis
fondation