



Robin noir N



Name of the variety in France

Robin noir

Origin

This variety is native to the Alpine vineyards. Based on genetic analyses carried out in Montpellier, Robin noir would result from the crossbreeding of Mondeuse blanche and Tressot.

Synonyms

In France, this variety can officially be called "Salagnin" regarding plant propagation material.

Legal information

In France, Robin noir is officially listed in the "Catalogue of vine varieties" since 2015 on the A list and classified.

Use

Wine grape variety.

Evolution of cultivated areas in France



Descriptive elements

The identification is based on:

the green tip of the young shoot with a low to medium density of prostrate hairs and a piping distribution of anthocyanin coloration,

- the shiny green young leaves with bronze spots and a very low to low density of prostrate hairs,
- the shoots with green or red-striped internodes,
- the wedge-shaped or circular adult leaves, entire or with three or five lobes, with moderately deep narrow upper lateral sinuses, an open V- or U-shaped petiole sinus, small to medium teeth, moderately long or long compared to their width at the base with straight sides or with one side concave and one side convex, no anthocyanin coloration of veins, a twisted, blistered, slightly gofferred leaf blade, sometimes folded near the petiole sinus, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,
- the ellipsoid or sometimes round-shaped berries.

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allel 1	131	223	249	176	194	256	238	216	239
Allel 2	137	229	260	186	200	260	240	257	261

Phenology

Bud burst: 5 days after Chasselas.

Grape maturity: early-season, 1 week and a half to 2 weeks after Chasselas.

Suitability for cultivation and agronomic production

Robin noir is vigorous, productive and has a semi-erect to erect bearing. This variety can sometimes be susceptible to millerandage.

Susceptibility to diseases and pests

Robin noir is very sensitive to downy mildew but it is not very affected by powdery mildew.

Technological potentiality

Robin noir's bunches are small to medium, often winged and compact. The berries are medium in size, simple-flavored, with a moderately thick skin and a juicy pulp. The berries have a good sugar accumulation potential but are not very acidic. This variety produces colored and tannic wines that are suited to ageing.

Clonal selection in France

There is no certified clone for this variety yet.

Bibliographic references

- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Montpellier SupAgro, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Registre Ampélographique International. 1961-1972, Ed. OIV, France.
- Le vignoble. A. Mas and V. Pulliat, 1874-1879, Ed. Masson, Paris, France.



Cette œuvre est mise à disposition selon les termes de la [Licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Partage dans les Mêmes Conditions 4.0 International](https://creativecommons.org/licenses/by-nc-sa/4.0/)

